

A Model of Blood Flow in a Circulation Network

Weihua Ruan[†], M.E. Clark[‡], Meide Zhao[‡] and Anthony Curcio[‡]

[†]Department of Mathematics, Computer Science and Statistics,
Purdue University Calumet

and

[‡]VasSol, Inc.

Abstract. We study a mathematical model of a blood circulation network which is a generalization of the coronary model proposed by Smith, Pullan and Hunter. We prove the existence and uniqueness of the solution to the initial-boundary value problem and discuss the continuity of dependence of the solution and its derivatives on initial, boundary and forcing functions and their derivatives.

1 Introduction

In a recent paper [13], Smith, Pullan and Hunter propose a mathematical model of blood circulation in the coronary network, and conduct a numerical analysis. In their model, major vessels (with cross-sectional areas larger than a certain value) are treated as a connected one-dimensional network, and small vessels, such as arterioles, capillaries and venules, are treated as lumped elements which are connected to the network of vessels. The flow on vessels are assumed to be incompressible, homogeneous, Newtonian, and has a small Reynolds number. Thus, the mass balance equation and Navier-Stokes equation can be written to describe the pressure and the flow rate on vessels. Equations on lumped elements are written in analogy with the current and voltage in an electric circuit. The result is an initial-boundary value problem of a system of hyperbolic type partial differential equations coupled at junctions of the network. Although the result of the numerical analysis conducted in [13] matches closely with measured data, the well-posedness problem of the system of partial differential equations, that is, the existence, uniqueness and the continuous dependence on initial and boundary data of the solution, has not been established before. The main objective of this paper is to establish the well-posedness. We prove that the system is well-posed under certain natural conditions. This work is an extension of our earlier work [12] on a model of blood circulation in the brain. The main differences between the two models are that the network configuration in [12] is more complicated owing to the presence of Willis loops, but the coupling junction conditions in the model of [13] are more complicated due to the different formulation and the inclusion of the capillaries and veinal system. We combine both features in a more general system with the hope that our result will be useful in the

modelling of circulation systems of higher complexity, including the whole body circulation system.

Before stating our system, let us briefly describe the model in [13]. Let P_i and R_i represent the pressure and radius on the i -th vessel, respectively, and let V_i be the cross-sectional average of the axial component $v_{i,x}$ of the velocity on the i -th vessel. Assuming that the radial component $v_{i,r}$ of the velocity is small compared to the axial component $v_{i,x}$ of the velocity, one can write equations of mass balance

$$\frac{\partial R_i}{\partial t} + V_i \frac{\partial R_i}{\partial x} + \frac{R_i}{2} \frac{\partial V_i}{\partial x} = 0, \quad (1.1)$$

and momentum balance

$$\frac{\partial V_i}{\partial t} + 2(1 - \alpha_i) \frac{V_i}{R_i} \frac{\partial R_i}{\partial t} + \alpha_i V_i \frac{\partial V_i}{\partial x} + \frac{1}{\rho} \frac{\partial P_i}{\partial x} = \frac{2\nu}{R_i} \left[\frac{\partial v_{i,x}}{\partial r} \right]_{r=R_i}. \quad (1.2)$$

Here ν is the viscosity constant and

$$\alpha_i = \frac{1}{R_i^2 V_i^2} \int_0^{R_i} 2r v_{i,x}^2 dr$$

is the energy quantity. Taking into consideration of no-slip boundary condition ($v_{i,x} = 0$ if $r = R_i$), the viscous axisymmetry ($\partial v_{i,x} / \partial r = 0$ if $r = 0$), and the fact that V_i is the cross-sectional average of $v_{i,x}$, Smith, Pullan and Hunter propose the velocity profile

$$v_{i,x}(r, x) = \frac{\gamma_i + 2}{\gamma_i} V_i(x) \left[1 - \left(\frac{r}{R_i} \right)^{\gamma_i} \right]$$

where γ_i is a positive number. Using this profile and the mass balance condition (1.1), Eq. (1.2) becomes

$$\frac{\partial V_i}{\partial t} + (2\alpha_i - 1) V_i \frac{\partial V_i}{\partial x} + 2(\alpha_i - 1) \frac{V_i^2}{R_i} \frac{\partial R_i}{\partial x} + \frac{1}{\rho} \frac{\partial P_i}{\partial x} = -\frac{2\nu\alpha_i}{\alpha_i - 1} \frac{V_i}{R_i^2} \quad (1.3)$$

with

$$\alpha_i = \frac{\gamma_i + 2}{\gamma_i} \in (1, \infty).$$

The pressure P_i and the radius R_i are related by a function

$$P_i = P_i(x, R_i).$$

In [13], it is assumed that

$$P_i(x, R_i) = C \left[\left(\frac{R_i}{R_0} \right)^\beta - 1 \right]$$

where C , R_0 and β are constants. We do not make such an assumption, only assume that it is a differentiable function and

$$\frac{\partial P_i}{\partial R_i} > 0$$

for all x and R_i . Let

$$A_i = \pi R_i^2, \quad Q_i = A_i V_i$$

be the cross-section area and the flow rate, respectively. It can be shown that the system of equations (1.1) and (1.3) is equivalent to

$$\begin{aligned} \frac{\partial A_i}{\partial t} + \frac{\partial Q_i}{\partial x} &= 0, \\ \frac{\partial Q_i}{\partial t} + \alpha_i \frac{\partial}{\partial x} \left(\frac{Q_i^2}{A} \right) + \frac{A_i}{\rho} \frac{\partial P_i}{\partial x} &= -\frac{4\pi\nu\alpha_i}{\alpha_i - 1} \frac{Q_i}{A_i}. \end{aligned} \quad (1.4)$$

Also, by rescaling the spatial variable x , we may assume that each vessel is parameterized to $x \in (0, 1)$.

The system of differential equations are supplemented with the initial condition

$$P_i(x, 0) = P_i^I(x), \quad Q_i(x, 0) = Q_i^I(x) \quad (1.5)$$

and boundary conditions. Boundary conditions at each end of the vessel are given according to the type of the end. If it is an external end of the network, either the pressure

$$P_i = P_i^B(t) \quad (1.6)$$

or the flow rate

$$Q_i = Q_i^B(t) \quad (1.7)$$

is specified. If the end is a *branching junction*, a junction connecting several vessels, let $j_1, \dots, j_\nu$ and $j_{\nu+1}, \dots, j_\mu$ denote the incoming and outgoing vessels, respectively. One imposes the mass balance condition

$$\sum_{l=1}^{\nu} Q_{j_l}(1, t) = \sum_{l'=\nu+1}^{\mu} Q_{j_{l'}}(0, t), \quad (1.8)$$

and the momentum balance condition

$$\rho_{j_l} \frac{\partial Q_{j_l}}{\partial t} = A_{j_l} (P_{j_l} - P_{j_{unc}}), \quad \rho_{j_{l'}} \frac{\partial Q_{j_{l'}}}{\partial t} = A_{j_{l'}} (P_{j_{unc}} - P_{j_{l'}}) \quad (1.9)$$

for $l = 1, \dots, \nu$, $l' = \nu+1, \dots, \mu$, where ρ_i are small positive constants and $P_{j_{unc}}$ is the pressure at the junction. (In [13], every branching junction connects only three vessels, it is either a

bifurcation point of one artery into two smaller ones or a joining point of two veins into a bigger one. Our prescription allows more general configuration of the network, including the presence of Willis loops.) If the end is a *transitional junction*, which connects the vessel to a network of arterioles, capillaries and venules, we follow the so called *microcirculation* model proposed in [13, 14]. Generalizing from [13], arterioles or venules connected to the vessel j_l are represented by a lumped resistive element R_{j_l} . The capillary bed is also represented by a resistive element R_C . R_{j_l} 's are connected to R_C through capacitive elements C_1 and C_2 on the two ends.




Figure 1: Microcirculation model of the network of arterioles, capillaries and venules.

Let $j_1, \dots, j_\nu$ be the arteries and let $j_{\nu+1}, \dots, j_\mu$ be the veins that are connected to a arteriole-capillary-venule network. The boundary conditions for P_{j_l} , Q_{j_l} are

$$\begin{aligned} R_{j_l} Q_{j_l}(1, t) &= P_{j_l}(1, t) - P_{C_1}(t), \quad \text{for } l = 1, \dots, \nu, \\ R_{j_{l'}} Q_{j_{l'}}(0, t) &= P_{C_2}(0, t) - P_{j_{l'}}(t), \quad \text{for } l' = \nu + 1, \dots, \mu \end{aligned} \quad (1.10)$$

and

$$C_1 \frac{dP_{C_1}}{dt} = \sum_{l=1}^{\nu} Q_{j_l}(1, t) - Q_C, \quad C_2 \frac{dP_{C_2}}{dt} = Q_C(t) - \sum_{l'=\nu+1}^{\mu} Q_{j_{l'}}(0, t) \quad (1.11)$$

where P_{C_i} , $i = 1, 2$ represent the pressure in the capacitive elements C_1 , C_2 , and

$$Q_C = \frac{P_{C_1} - P_{C_2}}{R_C} \quad (1.12)$$

represents the flow rate in the resistive element R_C . (In [13], there is only one artery and one vein connected to the system of arteriole-capillary-venule at the two ends. We do not rule out the possibility of multiple arteries and veins join together to such a system.)

The system we study in this paper consists of the equations

$$\begin{aligned} \frac{\partial P_i}{\partial t} + a_i \frac{\partial Q_i}{\partial x} &= f_i, \\ \frac{\partial Q_i}{\partial t} + b_i \frac{\partial P_i}{\partial x} + 2c_i \frac{\partial Q_i}{\partial x} &= g_i, \end{aligned} \quad x \in (0, 1), \quad t > 0 \quad (1.13)$$

and the initial and boundary conditions given by (1.5)–(1.12). For convenience, we also use the vector form

$$(U_i)_t + B_i (U_i)_x = F_i \quad (1.14)$$

where $U_i = (P_i, Q_i)$, $F_i = (f_i, g_i)$ and

$$B_i = \begin{pmatrix} 0 & a_i \\ b_i & 2c_i \end{pmatrix}.$$

Eq. (1.4) is a special case of this system where

$$a_i = \frac{\partial P_i}{\partial A_i}, \quad b_i = \frac{A_i}{\rho} - \frac{\alpha Q_i^2}{A_i^2} \left(\frac{\partial P_i}{\partial A_i} \right)^{-1}, \quad c_i = \frac{\alpha Q_i}{A_i}, \quad f_i = 0, \quad g_i = \frac{\alpha Q_i^2}{A_i^2} \frac{\partial A_i}{\partial x} - \frac{4\pi\nu\alpha}{\alpha - 1} \frac{Q_i}{A_i}.$$

We do not assume any particular form of these functions though, they are general differentiable functions of (x, t, P_i, Q_i) . Our basic assumptions are $a_i > 0$ and $A_i > \varepsilon_0$ for some positive constant ε_0 . Other assumptions will follow. Apart from the junction conditions, this system is the same as the one we study in [12]. Also, the junction conditions in [12] is the special case of (1.8)–(1.9) above with $\rho_i = 0$. As in [12], we use a fixed point principle to prove the solvability of the problem. Substituting a pair of functions (p_i, q_i) for (P_i, Q_i) in the coefficients a_i, b_i, c_i, A_i and forcing functions f_i, g_i , the system becomes linear. That is, all the functions a_i , etc. are independent of unknowns. If the linear system has a unique solution, then, one can establish a mapping from (p_i, q_i) to the linear problem solution (P_i, Q_i) . If one also shows that this mapping has a unique fixed point, then the fixed point is necessarily the unique solution of the quasilinear system. Hence, we shall first give a condition for the linear system to have a unique solution, then examine under what conditions the mapping has a unique fixed point. The first aspect of the problem is investigated in Section 2 and the second in Section 3. We also prove a result on the continuity of dependence of solutions on the initial, boundary and forcing functions for linear and quasilinear systems, thus, completing the analysis of the well-posedness of the problem. In spite of similarity in parts of the analysis to the one used in [12], the more general branching junction condition and the new transitional junction conditions require more careful treatments. Hence, there are substantial variations in the analysis. For completeness and to benefit the reader, we include all the major arguments in this paper.

2 The linear system

In this section, we analyze (1.13) as a linear system with $a_i, b_i, c_i, f_i, g_i, A_i$ independent of P_i and Q_i . The initial and boundary conditions are given by (1.5)–(1.12) except that the junction condition (1.9) is substituted by the more general condition

$$\rho_{j_l} \frac{\partial Q_{j_l}}{\partial t} = A_{j_l} (P_{j_l} - P_{j_{unc}}) + C_{j_l}, \quad \rho_{j_{l'}} \frac{\partial Q_{j_{l'}}}{\partial t} = A_{j_{l'}} (P_{j_{unc}} - P_{j_{l'}}) + C_{j_{l'}} \quad (2.1)$$

where C_i are differentiable functions of (x, t) . The inclusion of C_i is needed in the next section in order that the result of this section can be extended to the quasilinear system. We give conditions for the linear system to have a unique global solution. The conditions are most naturally given in terms of the eigenvalues of the matrix B_i , which have the form

$$\lambda_i^R = c_i + u_i, \quad \lambda_i^L = c_i - u_i,$$

where

$$u_i = \sqrt{c_i^2 + a_i b_i}.$$

These eigenvalues are real if

$$c_i^2 + a_i b_i > 0, \quad x \in (0, 1), \quad t > 0, \quad i = 1, \dots, n. \quad (2.2)$$

In this case,

$$\lambda_i^R(x, t) > 0, \quad \lambda_i^L(x, t) < \lambda_i^R(x, t) \quad (2.3)$$

and the system is hyperbolic. Under the condition (2.2), we show that the linear system has a unique solution if

$$\lambda_i^L(0, t) < 0, \quad \lambda_i^L(1, t) < 0, \quad i = 1, \dots, n$$

which is equivalent to

$$a_i b_i > 0, \quad t \geq 0, \quad i = 1, \dots, n. \quad (2.4)$$

at $x = 0, 1$ only. It needs not hold for $x \in (0, 1)$.

Theorem 2.1 *Assume that the functions $a_i, b_i, c_i, f_i, g_i, A_i$ and C_i are independent of (P_i, Q_i) . Suppose that these functions and the initial and boundary functions $P_i^I, Q_i^I, P_i^B, Q_i^B$ all have bounded first-order derivatives. Suppose also that $a_i > 0, A_i > 0$ and that the conditions (2.2) and (2.4) hold. Then, for any $T > 0$ there is a unique solution in a bounded subset of the space $C([0, 1] \times [0, T], \mathbb{R}^{2n})$ to the linear system (1.13) with the initial and boundary conditions given by (1.5)–(1.8), (1.10)–(1.12), and (2.1).*

Proof. We first show that the system has a unique solution for $0 < t < \delta$ for some $\delta > 0$. The proof is based on the method of characteristics and a fixed point principle. For systems defined on only one branch with boundary conditions of the forms of (1.6) or (1.7), this is a standard approach. In our case, special care is needed to handle the junction conditions.

Consider the i -th branch. From any point (ξ, τ) on the left, right, and lower boundary of the rectangle $D =: [0, 1] \times [0, T]$, we construct the left-going and right-going characteristic curves $x = x_i^L(t; \xi, \tau)$ and $x = x_i^R(t; \xi, \tau)$ by

$$\begin{aligned} \frac{dx_i^L}{dt} &= \lambda_i^L(x_i^L, t), \quad x_i^L(\tau) = \xi, \\ \frac{dx_i^R}{dt} &= \lambda_i^R(x_i^R, t), \quad x_i^R(\tau) = \xi, \end{aligned}$$

respectively, where λ_i^L and λ_i^R are the two eigenvalues of the matrix B_i . By the uniqueness of solutions to these differential equations, a left-going (resp. right-going) characteristic curve cannot intersect with another left-going (resp. right-going) characteristic curve. Let X_i^L and X_i^R be the right-most left-going and left-most right-going characteristic curves,

$$x = x_i^L(t; 1, 0) \quad \text{and} \quad x = x_i^R(t; 0, 0)$$

starting from the lower boundary of D , respectively. It can be shown from (2.3) that the two curves can have at most one intersection. Let t_i be the value of t at the intersection. If the two curves do not intersect in D , we simply define $t_i = T$. By condition (2.4), X_i^L cannot reach the right vertical line $x = 1$ at any $t > 0$, and by $\lambda_i^R > 0$, X_i^R cannot reach the vertical line $x = 0$ at any $t > 0$. Thus, the rectangle $D_i =: [0, 1] \times [0, t_i]$ can be divided into three parts

$$D_i = D_i^L \cup D_i^C \cup D_i^R,$$

where D_i^L is between the vertical line $x = 0$ and the characteristic curve X_i^R , D_i^C is between the two characteristic curves, and D_i^R is between X_i^L and $x = 1$.




Figure 2: Three parts of D_i

We show that there is a $\delta_i \leq t_i$ such that the solution (P_i, Q_i) for the i -th branch exists in the restriction of D_i to the strip $\{0 \leq t \leq \delta_i\}$.

First, observe that the initial conditions alone determine the solution completely in the central region D_i^C . This follows from the theory of first-order linear hyperbolic systems and the fact that from any point $(x, t) \in D_i^C$, the two characteristic curves, followed backwards, must land on the horizontal line $t = 0$. (The latter is a consequence of (2.3).) To extend the solution to other parts of D_i , we make a change of unknowns and derive a set of integral equations. Note that $l_i^R =: (-\lambda_i^L, a_i)$ and $l_i^L =: (-\lambda_i^R, a_i)$ are the left eigenvectors of B_i corresponding to λ_i^R and λ_i^L , respectively. Introduce new unknowns

$$r_i = l_i^R U_i \equiv -\lambda_i^L P_i + a_i Q_i, \quad s_i = l_i^L U_i \equiv -\lambda_i^R P_i + a_i Q_i. \quad (2.5)$$

The system (1.13) can be written in terms of r_i and s_i by multiplying the left eigenvectors to (1.14) and substituting in

$$P_i = \frac{1}{2u_i} (r_i - s_i), \quad Q_i = \frac{1}{2u_i a_i} (\lambda_i^R r_i - \lambda_i^L s_i). \quad (2.6)$$

This results in the equations

$$\partial_i^R r_i = F_i^R(x, t, r_i, s_i), \quad \partial_i^L s_i = F_i^L(x, t, r_i, s_i), \quad (2.7)$$

where

$$\partial_i^R = \frac{\partial}{\partial t} + \lambda_i^R \frac{\partial}{\partial x}, \quad \partial_i^L = \frac{\partial}{\partial t} + \lambda_i^L \frac{\partial}{\partial x}, \quad (2.8)$$

and

$$F_i^R(x, t, r_i, s_i) = l_i^R F_i + (\partial_i^R l_i^R) U_i, \quad F_i^L(x, t, r_i, s_i) = l_i^L F_i + (\partial_i^L l_i^L) U_i. \quad (2.9)$$

(A differential operator acting on a vector means that it acts on each component of the vector.) Let $(x, t) \in D_i$. We integrate the first equation of (2.7) along the right-going characteristic curve $x^R(t; \xi, \tau)$ which passes through (x, t) and reaches the left or lower boundary of D_i at (ξ, τ) . It can be shown that for $(x, t) \in D_i^C \cup D_i^R$, $\tau = 0$, and for $(x, t) \in D_i^L$, $\xi = 0$. In the former case, we obtain

$$r_i(x, t) = r_i^I(\xi) + \int_0^t F_i^R(x_i^R(t'; \xi, 0), t', r_i, s_i) dt' \quad (2.10)$$

In the latter case, we have

$$r_i(x, t) = r_i(0, \tau) + \int_\tau^t F_i^R(x_i^R(t'; 0, \tau), t', r_i, s_i) dt'. \quad (2.11)$$

Similarly, by integrating the second equation of (2.7) along the left-going characteristic curve $x_i^L(t; \xi, \tau)$ that passes through both (x, t) and (ξ, τ) (which is on either the right or lower boundary of D_i), the equations are

$$s_i(x, t) = s_i^I(\xi) + \int_0^t F_i^L(x_i^L(t'; \xi, 0), t', r_i, s_i) dt' \quad (2.12)$$

if $(x, t) \in D_i^L \cup D_i^C$ and

$$s_i(x, t) = s_i(1, \tau) + \int_\tau^t F_i^L(x_i^L(t'; 1, \tau), t', r_i, s_i) dt' \quad (2.13)$$

if $(x, t) \in D_i^R$. These are the integral equations we need.

For any $\delta_i \leq t_i$ we use D_{i, δ_i}^L , D_{i, δ_i}^C and D_{i, δ_i}^R to denote the restrictions of D_i^L , D_i^C and D_i^R to the strip $\{0 \leq t \leq \delta_i\}$, respectively. First, consider the case where the end of the branch is an external end. We discuss the case of a left end only, the case of a right end can be treated similarly. If the boundary condition is given by (1.6), we define $\hat{s}_i = s_i/\varepsilon$ where $\varepsilon < 1$ is any constant. Using the first equation of (2.6) in the integral equations (2.11) and (2.12),

$$\begin{pmatrix} r_i(x, t) \\ \hat{s}_i(x, t) \end{pmatrix} = \begin{pmatrix} 2u_i(0, \tau) P_i^B(\tau) + \varepsilon \hat{s}_i(0, \tau) + \int_\tau^t F_i^R(x_i^R(t'; 0, \tau), t', r_i, \varepsilon \hat{s}_i) dt' \\ \frac{1}{\varepsilon} s_i^I(\xi) + \frac{1}{\varepsilon} \int_0^t F_i^L(x_i^L(t'; \xi, 0), t', r_i, \varepsilon \hat{s}_i) dt' \end{pmatrix}. \quad (2.14)$$

This is a fixed point equation for $(r_i, \hat{s}_i)$ if we define the right hand side as a mapping of an operator K on $(r_i, \hat{s}_i)$ in a bounded subset of $C(D_{i, \delta_i}^L \cup D_{i, \delta_i}^C, \mathbb{R}^2)$. In a standard approach, it can be shown that K is a contraction mapping if δ_i is sufficiently small. Hence, the fixed point exists and is unique, and the solution (r_i, s_i) is uniquely extended to $D_{i, \delta_i}^L \cup D_{i, \delta_i}^C$. If the boundary condition is given by (1.7), we define $\hat{s}_i = s_i/\varepsilon$, where $\varepsilon > 0$ is so small such that

$$\varepsilon \left| \frac{\lambda_i^L(0, \tau)}{\lambda_i^R(0, \tau)} \right| < 1, \quad \tau \in (0, t_i).$$

The fixed point equation is then

$$\begin{pmatrix} r_i(x, t) \\ \hat{s}_i(x, t) \end{pmatrix} = \begin{pmatrix} \frac{2a_i u_i(0, \tau)}{\lambda_i^R(0, \tau)} Q_i^B(\tau) + \frac{\lambda_i^L(0, \tau)}{\lambda_i^R(0, \tau)} \varepsilon \hat{s}_i(0, \tau) + \int_\tau^t F_i^R(x_i^R(t'; 0, \tau), t', r_i, \varepsilon \hat{s}_i) dt' \\ \frac{1}{\varepsilon} s_i^I(\xi) + \frac{1}{\varepsilon} \int_0^t F_i^L(x_i^L(t'; \xi, 0), t', r_i, \varepsilon \hat{s}_i) dt' \end{pmatrix}. \quad (2.15)$$

By a similar argument, the solution can again be uniquely extended.

We next extend the solution to either D_{i, δ_i}^L or D_{i, δ_i}^R if the end is a branching junction. In this case, we shall extend the solution on all the branches that are connected to the

same junction simultaneously. Let $j_1, \dots, j_\nu$ be the incoming and $j_{\nu+1}, \dots, j_\mu$ the outgoing branches to the junction. Equations (1.8), (2.1) and (2.6) give rise to a $2\mu \times \mu$ homogenous system of linear (ordinary) differential equations for $r_i(1, t)$, $s_i(1, t)$, $i = j_1, \dots, j_\nu$ and $r_i(0, t)$, $s_i(0, t)$, $i = j_{\nu+1}, \dots, j_\mu$:

$$\begin{aligned} \frac{\rho_{j_1}}{A_{j_1}} \frac{d}{dt} Q_{j_1}(1, t) - \frac{C_{j_1}}{A_{j_1}} - P_{j_1}(1, t) &= \frac{\rho_i}{A_i} \frac{d}{dt} Q_i(1, t) - \frac{C_i}{A_i} - P_i(1, t), \quad i = j_2, \dots, j_\nu, \\ \frac{\rho_{j_1}}{A_{j_1}} \frac{d}{dt} Q_{j_1}(1, t) - \frac{C_{j_1}}{A_{j_1}} - P_{j_1}(1, t) &= -\frac{\rho_i}{A_i} \frac{d}{dt} Q_i(0, t) - \frac{C_i}{A_i} - P_i(0, t), \quad i = j_{\nu+1}, \dots, j_\mu, \\ \sum_{l=1}^{\nu} Q_{j_l}(1, t) - \sum_{l'=\nu+1}^{\mu} Q_{j_{l'}}(0, \tau) &= 0. \end{aligned} \tag{2.16}$$

Differentiate the last equation with respect to t and regard $s_{j_1}(1, t), \dots, s_{j_\nu}(1, t), r_{j_{\nu+1}}(0, t), \dots, r_{j_\mu}(0, t)$ as unknowns. The derivatives of unknowns can be solved from (2.16) because the coefficient matrix of ds_i/dt and dr_i/dt in (2.16),

$$\begin{pmatrix} -\frac{\rho_{j_1} \lambda_{j_1}^L(1, t)}{2u_{j_1} a_{j_1} A_{j_1}(1, t)} & \frac{\rho_{j_2} \lambda_{j_2}^L(1, t)}{2u_{j_2} a_{j_2} A_{j_2}(1, t)} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ -\frac{\rho_{j_1} \lambda_{j_1}^L(1, t)}{2u_{j_1} a_{j_1} A_{j_1}(1, t)} & 0 & \dots & \frac{\rho_{j_\mu} \lambda_{j_\mu}^R(0, t)}{2u_{j_\mu} a_{j_\mu} A_{j_\mu}(0, t)} \\ -\frac{\lambda_{j_1}^L(1, t)}{2u_{j_1} a_{j_1}(1, t)} & -\frac{\lambda_{j_2}^L(1, t)}{2u_{j_2} a_{j_2}(1, t)} & \dots & -\frac{\lambda_{j_\mu}^R(0, t)}{2u_{j_\mu} a_{j_\mu}(0, t)} \end{pmatrix}$$

has the determinant

$$\left(\frac{-1}{2}\right)^\mu \prod_{l=1}^{\nu} \frac{\rho_{j_l} \lambda_{j_l}^L(1, t)}{u_{j_l} a_{j_l} A_{j_l}(1, t)} \prod_{l'=\nu+1}^{\mu} \frac{\rho_{j_{l'}} \lambda_{j_{l'}}^R(0, t)}{u_{j_{l'}} a_{j_{l'}} A_{j_{l'}}(0, t)} \sum_{l=1}^{\mu} \frac{A_{j_l}}{\rho_{j_l}}.$$

Since $\lambda_i^L < 0 < \lambda_i^R$ at the junction, the determinant is not zero. Thus, the derivatives of the unknowns, $s_{j_l}(1, t)$ and $r_{j_{l'}}(0, t)$, are each a linear combination of the functions $r_{j_l}(1, t)$, $s_{j_l}(1, t)$, $r_{j_{l'}}(0, t)$, $s_{j_{l'}}(0, t)$ together with the derivatives of $r_{j_l}(1, t)$ and $s_{j_{l'}}(0, t)$, $l = 1, \dots, \nu$, $l' = \nu + 1, \dots, \mu$. Integrating and using the initial condition determined by (1.5) and (2.5), we can write

$$s_i(1, \tau) = s_i(1, 0) + \sum_{l=1}^{\nu} m_{j_l}^i(\tau) r_{j_l}(1, \tau) + \sum_{l'=\nu+1}^{\mu} m_{j_{l'}}^i(\tau) s_{j_{l'}}(0, \tau) + \int_0^\tau H_i dt', \tag{2.17}$$

for $i = j_1, \dots, j_\nu$ and

$$r_i(0, \tau) = r_i(0, 0) + \sum_{l=1}^{\nu} n_{j_l}^i(\tau) r_{j_l}(1, \tau) + \sum_{l'=\nu+1}^{\mu} n_{j_{l'}}^i(\tau) s_{j_{l'}}(0, \tau) + \int_0^\tau H_i dt', \tag{2.18}$$

for $i = j_{\nu+1}, \dots, j_\mu$, where m_j^i, n_j^i are continuous functions and H_i are linear combinations of $r_{j_l}(1, t), s_{j_l}(1, t), r_{j_{l'}}(0, t), s_{j_{l'}}(0, t)$ and $C_i(t)$ with coefficients depending only on t . Choose an $\varepsilon > 0$ such that

$$\varepsilon \max \left\{ \sum_{l=1}^{\mu} |m_{j_l}^i(\tau)|, \sum_{l=1}^{\mu} |n_{j_l}^i(\tau)| \right\} < 1, \quad i = j_1, \dots, j_\mu, \quad \tau \in [0, t_i]$$

and introduce

$$\hat{r}_{j_l} = \frac{r_{j_l}}{\varepsilon}, \quad \hat{s}_{j_{l'}} = \frac{s_{j_{l'}}}{\varepsilon}, \quad l = 1, \dots, \nu, \quad l' = \nu + 1, \dots, \mu.$$

Then, from (2.10)–(2.13), the integral equations for the 2μ unknowns $\hat{r}_{j_l}, s_{j_l}, r_{j_{l'}}, \hat{s}_{j_{l'}}$, $l = 1, \dots, \nu, l' = \nu + 1, \dots, \mu$ constitute a fixed point equation, $w = Kw$, where

$$w = (\hat{r}_{j_1}, \dots, \hat{r}_{j_\nu}, s_{j_1}, \dots, s_{j_\nu}, r_{j_{\nu+1}}, \dots, r_{j_\mu}, \hat{s}_{j_{\nu+1}}, \dots, \hat{s}_{j_\mu}) \quad (2.19)$$

and

$$\begin{aligned} Kw = & \left(\frac{1}{\varepsilon} r_{j_1}^I(\xi_{j_1}) + \frac{1}{\varepsilon} \int_0^t F_{j_1}^R dt', \dots, \right. \\ & s_{j_1}(1, 0) + \varepsilon \left(\sum_{k=1}^{\nu} m_{j_k}^1 \hat{r}_{j_k}(1, \tau) + \sum_{k'=\nu+1}^{\mu} m_{j_{k'}}^1 \hat{s}_{j_{k'}}(0, \tau) \right) + \int_0^\tau H_{j_1} dt' + \int_\tau^t F_{j_1}^L dt', \dots, \\ & r_{j_{\nu+1}}(0, 0) + \varepsilon \left(\sum_{k=1}^{\nu} n_{j_k}^1 \hat{r}_{j_k}(1, \tau) + \sum_{k'=\nu+1}^{\mu} n_{j_{k'}}^1 \hat{s}_{j_{k'}}(1, \tau) \right) + \int_0^\tau H_{j_{\nu+1}} dt' + \int_\tau^t F_{j_{\nu+1}}^R dt', \dots, \\ & \left. \frac{1}{\varepsilon} s_{j_{\nu+1}}^I(\xi_{j_{\nu+1}}) + \frac{1}{\varepsilon} \int_0^t F_{j_{\nu+1}}^L dt', \dots \right) \end{aligned} \quad (2.20)$$

in which

$$F_{j_l}^R = F_{j_l}^R(x_{j_l}^R, t', \varepsilon \hat{r}_{j_l}, s_{j_l}), \quad F_{j_l}^L = F_{j_l}^L(x_{j_l}^L, t', \varepsilon \hat{r}_{j_l}, s_{j_l})$$

for $l = 1, \dots, \nu$, and

$$F_{j_{l'}}^R = F_{j_{l'}}^R(x_{j_{l'}}^R, t', r_{j_{l'}}, \varepsilon \hat{s}_{j_{l'}}), \quad F_{j_{l'}}^L = F_{j_{l'}}^L(x_{j_{l'}}^L, t', r_{j_{l'}}, \varepsilon \hat{s}_{j_{l'}})$$

for $l' = \nu + 1, \dots, \mu$. It can be shown by a standard argument that K is a contraction mapping in the space

$$X_j =: \prod_{l=1}^{\nu} C(D_{j_l, \delta_j}^C \cup D_{j_l, \delta_j}^R, \mathbb{R}^2) \times \prod_{l=\nu+1}^{\mu} C(D_{j_l, \delta_j}^L \cup D_{j_l, \delta_j}^L, \mathbb{R}^2)$$

if δ_j is sufficiently small. Hence, it has a unique fixed point in X_j . This extends the solution (r_i, s_i) for the neighboring branches of the junction.

It remains to extend the solution to a region adjacent to a transitional junction. Similar to the case of a branching junction, we simultaneously treat all the branches that are connected to the same transitional junction. Let $j_1, \dots, j_\nu$ be the arteries and $j_{\nu+1}, \dots, j_\mu$ be the veins.

The condition connecting the vessels are given by (1.10), (1.11) and (1.12). Differentiate the equations in (1.10) with respect to t , the resulting equations together with (1.11) is a linear system of the derivatives of the functions $r_{j_l}(1, t)$, $s_{j_l}(1, t)$, $r_{j_{l'}}(0, t)$, $s_{j_{l'}}(0, t)$, $l = 1, \dots, \nu$, $l' = \nu + 1, \dots, \mu$, and $P_{C_1}(t)$, $P_{C_2}(t)$. The coefficient matrix of $ds_{j_l}(1, t)/dt$, $dr_{j_{l'}}(0, t)/dt$, for $l = 1, \dots, \nu$, $l' = \nu + 1, \dots, \mu$ and P'_{C_1} , P'_{C_2} is

$$\begin{bmatrix} D_1 & 0 & B_1 \\ 0 & D_2 & B_2 \\ 0 & 0 & I_2 \end{bmatrix}$$

where

$$\begin{aligned} D_1 &= \text{diag} \left(-\frac{R_{j_1} \lambda_{j_1}^L}{2u_{j_1} a_{j_1}} + \frac{1}{2u_{j_1}}, \dots, -\frac{R_{j_\nu} \lambda_{j_\nu}^L}{2u_{j_\nu} a_{j_\nu}} + \frac{1}{2u_{j_\nu}} \right), \\ D_2 &= \text{diag} \left(\frac{R_{j_{\nu+1}} \lambda_{j_{\nu+1}}^R}{2u_{j_{\nu+1}} a_{j_{\nu+1}}} + \frac{1}{2u_{j_{\nu+1}}}, \dots, \frac{R_{j_\mu} \lambda_{j_\mu}^R}{2u_{j_\mu} a_{j_\mu}} + \frac{1}{2u_{j_\mu}} \right), \end{aligned}$$

I_2 is the 2×2 identity matrix, and B_1 , B_2 are some constant matrices. Since all the elements of the diagonal matrices D_1 and D_2 are positive, the system can be uniquely solved for these derivatives. Thus, each of $ds_{j_l}(1, t)/dt$, $dr_{j_{l'}}(0, t)/dt$ for $l = 1, \dots, \nu$, $l' = \nu + 1, \dots, \mu$ is a linear combination of $r_{j_l}(1, t)$, $s_{j_l}(1, t)$, $r_{j_{l'}}(0, t)$, $s_{j_{l'}}(0, t)$, and $dr_{j_l}(1, t)/dt$, $ds_{j_{l'}}(0, t)/dt$, $l = 1, \dots, \nu$, $l' = \nu + 1, \dots, \mu$ as well as P_{C_1} and P_{C_2} . We can also eliminate P_{C_1} and P_{C_2} by (1.10). Integrating the resulting equations, we obtain equations (2.17)–(2.18) for some functions m_j^i , n_j^i , and H_i . The remaining of the previous paragraph can then be used here to give the extension of the solution to the left or right regions for the branches.

Finally, if we let δ be the minimum of all δ_i occurring above, we see that $\delta > 0$ and the solution exists and is unique in $(x, t) \in D_\delta =: [0, 1] \times [0, \delta]$. Observe that δ depends only on the bounds of the system functions a_i , etc., the initial and boundary functions P_i^I , etc., and their first-order derivatives in $D = [0, 1] \times [0, T]$. Hence, it is independent of t , and we can extend the solution successively in the time intervals $[0, \delta]$, $[\delta, 2\delta]$, etc. In this way, the solution is obtained in D in finitely many steps. ■

We next derive an estimate of the deviation of solution in term of the deviations of the initial, boundary and forcing functions. This estimate is needed in the next section. For any vector function $v = (v_1, \dots, v_k)$ defined in $C(X; \mathbb{R}^k)$, we use $|v|_X$ to denote the norm $\max_i \left\{ |v_i|_{C(X)} \right\}$, where X represents a closed subset of either $\mathbb{R}$ or $\mathbb{R}^2$.

Lemma 2.1 *Let $U = (P, Q)$ and $\tilde{U} = (\tilde{P}, \tilde{Q})$ be two solutions of the linear problem (1.14) with different initial, boundary, and forcing functions. Suppose the conditions of Theorem 2.1 hold for both solutions. Suppose also that there is a positive lower bound for all A_i . Then,*

there exists a constant $M > 0$, independent of initial, boundary and forcing functions, such that

$$\begin{aligned} |U - \tilde{U}|_{C(D_\delta)} \leq M \Big(& |P^I - \tilde{P}^I|_{C[0,1]} + |Q^I - \tilde{Q}^I|_{C[0,1]} + |P^B - \tilde{P}^B|_{C[0,\delta]} + |Q^B - \tilde{Q}^B|_{C[0,\delta]} \\ & + |C - \tilde{C}|_{C[0,\delta]} + \delta |f - \tilde{f}|_{C(D_\delta)} + \delta |g - \tilde{g}|_{C(D_\delta)} \Big). \end{aligned} \quad (2.21)$$

Proof. We need only prove (2.21) for a $\delta \leq \min_i \{\delta_i\}$, where δ_i represents the constants occurring in the proof of Theorem 2.1. This is because for larger δ , we can divide the interval $[0, \delta]$ into subintervals, each has a length less than $\min_i \{\delta_i\}$, and apply (2.21) in each subinterval. We can then take the maximum on each side of the inequalities to derive the inequality of in $[0, \delta]$. In the sequel, D_δ^C , D_δ^L and D_δ^R are the restrictions of D_i^C , D_i^L and D_i^R to the strip $\{0 \leq t \leq \delta\}$, respectively.

By linearity, $U - \tilde{U}$ is the solution of the system with the initial, boundary and forcing functions $P_i^I - \tilde{P}_i^I$, $Q_i^I - \tilde{Q}_i^I$, $P_i^B - \tilde{P}_i^B$, $Q_i^B - \tilde{Q}_i^B$, $f_i - \tilde{f}_i$, $g_i - \tilde{g}_i$ and $C_i - \tilde{C}_i$. Let r_i , $\hat{r}_i$, s_i , $\hat{s}_i$ be defined as in the proof of Theorem 2.1, corresponding to $U - \tilde{U}$. We show that these quantities have upper bounds in the form of the right hand side of (2.21) in D_δ^C , D_δ^L and D_δ^R .

In D_δ^C , (2.10) and (2.12) hold. Notice that the functions F_i^R and F_i^L are linear in r_i , and s_i . Hence, there exists a constant M (we will use M generically for any constant bounds that are independent of solutions) such that

$$R_i^C(t) + S_i^C(t) \leq |r_i^I|_{C[0,1]} + |s_i^I|_{C[0,1]} + M \int_0^t (R_i^C(t') + S_i^C(t') + T_i^C(t')) dt',$$

where

$$R_i^C(t) = \sup_{\{x:(x,t) \in D_\delta^C\}} |r_i(x,t)|, \quad S_i^C(t) = \sup_{\{x:(x,t) \in D_\delta^C\}} |s_i(x,t)|, \quad (2.22)$$

and

$$T_i^C(t) = \sup_{\{x:(x,t) \in D_\delta^C\}} \left(|f_i(x,t) - \tilde{f}_i(x,t)| + |g_i(x,t) - \tilde{g}_i(x,t)| \right). \quad (2.23)$$

Hence, by Gronwall's inequality (see, e.g. [8, p.327]),

$$R_i^C(t) + S_i^C(t) \leq M \left(|r_i^I|_{C[0,1]} + |s_i^I|_{C[0,1]} + \delta \sup_{t \in (0,\delta)} T_i^C(t) \right)$$

for $t \in [0, \delta]$. This proves that R_i^C and S_i^C have upper bounds in the form of the right side of (2.21).

We next consider the left or right regions if it is adjacent to an external end. Since the both cases are similar, we will only treat the case where the left end is an external. The integral equations to be used are (2.14) or (2.15) according to the type of the boundary condition. The resulting inequality has the form

$$R_i^L(t) + \hat{S}_i^L(t) \leq \sigma \hat{S}_i^L(t) + M \left(|s_i^I|_{C[0,1]} + |\xi_i^B|_{C[0,\delta]} + \int_0^t \left(R_i^L(\tau) + \hat{S}_i^L(\tau) + T_i^L(\tau) \right) d\tau \right)$$

where ξ_i^B is either P_i^B or Q_i^B depending on the boundary condition, and R_i^L , $\hat{S}_i^L$ and T_i^L are defined in the same way as in (2.22)–(2.23), with D_δ^C substituted by $D_\delta^L \cup D_\delta^C$, and $\sigma > 0$ is a positive constant such that $\sigma = \varepsilon$ if the boundary condition is (1.6) and

$$\sigma = \varepsilon \sup_{t \in (0,\delta)} \left| \frac{\lambda_i^L(0,t)}{\lambda_i^R(0,t)} \right| < 1$$

if the boundary condition is (1.7). Replacing M by $(1 - \sigma)M$, we can write

$$R_i^L(t) + \hat{S}_i^L(t) \leq M \left(|s_i^I|_{C[0,1]} + |\xi_i^B|_{C[0,\delta]} + \int_0^t \left(R_i^L(\tau) + \hat{S}_i^L(\tau) + T_i^L(\tau) \right) d\tau \right).$$

Hence, by Gronwall's inequality

$$R_i^L(t) + \hat{S}_i^L(t) \leq M \left(|s_i^I|_{C[0,1]} + |\xi_i^B|_{C[0,\delta]} + \delta \max_{t \in (0,\delta)} T_i^L(t) \right).$$

This proves that both $R_i^L(t)$ and $\hat{S}_i^L(t)$ have upper bounds in the form of the right hand side of (2.21).

We next extend the estimate to $D_{i,\delta}^L$ or $D_{i,\delta}^R$ if the end is either a branching junction or a transitional junction. In either case, the solutions on the branches $j_1, \dots, j_\mu$ connecting to the junction constitute a fixed point of the operator K , which is defined in (2.20). Let

$$W(t) = \sum_{l=1}^{\nu} \left(\hat{R}_{j_l}^R(t) + S_{j_l}^R(t) \right) + \sum_{l'=\nu+1}^{\mu} \left(R_{j_{l'}}^L(t) + \hat{S}_{j_{l'}}^L(t) \right)$$

where $\hat{R}_i^R$ and S_i^R are defined as in (2.22) with D_δ^C substituted by $D_\delta^C \cup D_\delta^R$. Then, from $w = Kw$ and in view of the assumption that A_i has a positive lower bound for all i and $t > 0$, we can deduce

$$\begin{aligned} W(t) &\leq \sigma \left(\sum_{l=1}^{\nu} \hat{R}_{j_l}^R(t) + \sum_{l'=\nu+1}^{\mu} \hat{S}_{j_{l'}}^L(t) \right) \\ &\quad + M \left(\sum_{l=1}^{\nu} |r_{j_l}^I|_{C[0,1]} + \sum_{l'=\nu}^{\mu} |s_{j_{l'}}^I|_{C[0,1]} + \int_0^t (W(\tau) + T(\tau)) d\tau \right), \end{aligned}$$

where

$$T(\tau) = \sum_{l=1}^{\nu} T_{j_l}^R(\tau) + \sum_{l'=\nu+1}^{\mu} T_{j_{l'}}^L(\tau) + \sum_{l=1}^{\mu} \left| C^{j_l}(\tau) - \tilde{C}^{j_l}(\tau) \right|$$

and $T_i^R(t)$ is defined as in (2.23) with D_δ^C substituted by $D_\delta^C \cup D_\delta^R$. Replacing M by $(1 - \sigma)M$, we obtain

$$W(t) \leq M \left(\sum_{l=1}^{\nu} |r_{j_l}^I|_{C[0,1]} + \sum_{l'=\nu}^{\mu} |s_{j_{l'}}^I|_{C[0,1]} + \int_0^t (W(\tau) + T(\tau)) d\tau \right).$$

Hence, by Gronwall's inequality,

$$W(t) \leq M \left(\sum_{l=1}^{\nu} |r_{j_l}^I|_{C[0,1]} + \sum_{l'=\nu}^{\mu} |s_{j_{l'}}^I|_{C[0,1]} + \delta \max_{t \in (0, \delta)} T(t) \right).$$

This leads to an upper bound in the form of the right hand side of (2.21) for $R_i^R(t)$, $S_i^R(t)$, $i = j_1, \dots, j_\nu$, and $R_i^L(t)$, $S_i^L(t)$, $i = j_{\nu+1}, \dots, j_\mu$.

We have thus obtained an upper bound in the form of the right hand side of (2.21) for the quantities $|r_i - \tilde{r}_i|_{C(D_\delta)}$ and $|s_i - \tilde{s}_i|_{C(D_\delta)}$. The conclusion of the lemma follows now from (2.6). ■

3 The quasilinear system

In this section, we study the quasilinear system where the coefficients a_i , b_i , c_i , f_i , g_i , A_i and C_i depend on both (x, t) and (P_i, Q_i) . Under certain conditions, we show that the system has a unique local solution. We then present a theorem on the continuity of dependence of the solution on initial, boundary and forcing function.

The basic idea in the proof of the existence of solution is to construct an iterative sequence. Substituting any vector function (p_i, q_i) for (P_i, Q_i) in a_i , etc., the system becomes linear. Thus, we can use Theorem 2.1 to get a solution (P_i, Q_i) . This defines a mapping S from $u =: (p_i, q_i)$ to $U =: (P_i, Q_i)$, and the solution for the quasilinear system is a fixed point of S . If there is a subset of a Banach space that is invariant under S , then, we can construct a sequence

$$u_{k+1} = Su_k, \quad k = 0, 1, \dots$$

In the case where the limit exists and is unique, it gives rise to fixed point of S . This is our approach in this section.

In this approach, conditions (2.2) and (2.4) are repeatedly used. One might want to impose them for all the values of the variables. This would give the existence and uniqueness for the global solution, as in the case of the linear system. However, such a requirement is

so restrictive that even the original system (1.4) cannot meet it. Therefore, we will impose them only for $t = 0$, and obtain the local solution for the quasilinear system.

Theorem 3.1 *Assume that the initial and boundary functions $P_i^I, Q_i^I, P_i^B, Q_i^B$ and the system functions $a_i, b_i, c_i, f_i, g_i, A_i$ and C_i all have continuous first-order derivatives with respect to each variable. Suppose that a_i is positive and A_i has a positive lower bound for all the values of their arguments, and that conditions (2.2)–(2.4) hold at $t = 0$. Suppose also that the initial functions P_i^I, Q_i^I satisfy any relevant boundary conditions at $t = 0$. Then, for some $\delta > 0$, there is a unique solution for $0 \leq t < \delta$ to the quasilinear system (1.13) with the initial and boundary conditions given by (1.5)–(1.8), (2.1), and (1.10)–(1.12).*

Proof. We first consider the simpler case where $U^I =: (P^I, Q^I) = 0$. Let $v = \{v_i\}$, $v_i = (p_i, q_i)$ be a family of vector functions (not necessarily constitutes a solution) that satisfy the initial and boundary conditions. Substitute v for U in the functions $a_i, b_i, c_i, f_i, g_i, A_i$ and C_i . Then, the system becomes linear and we can invoke Theorem 2.1 to obtain a solution U to the linear system. This defines a mapping $S : v \mapsto U$. A solution to the quasilinear system is then a fixed point of S . We will choose a subset X_{δ, M_0} of a Banach space such that (1) $SX_{\delta, M_0} \subset X_{\delta, M_0}$, and (2) S is contracting in X_{δ, M_0} . For any scalar or vector function $f \in C^k(D_\delta)$, let $|f|_{k, \delta}$ denote the maximum norm of all the k -th order derivatives of f in D_δ . (If f is a vector function, $|f|_{k, \delta} = \max_i \{|f_i|_{k, \delta}\}$.) Let $C_B(D_\delta, \mathbb{R}^{2n})$ denote the subset of the vector-valued functions in $C(D_\delta, \mathbb{R}^{2n})$ that satisfy the initial and boundary conditions. We seek X_{δ, M_0} in the form

$$X_{\delta, M_0} = \left\{ v \in C_B(D_\delta, \mathbb{R}^{2n}) : |v|_{0, \delta} \leq M_0, |v|_{1, \delta} \leq M_1 \right\} \quad (3.1)$$

where M_0 is an arbitrary positive constant and M_1 is a constant to be determined. Note that by the vanishing initial condition, for any M_1 , $|U|_{1, \delta} \leq M_1$ implies $|U|_{0, \delta} \leq M_1 \delta$. Hence, for any M_0 , we can ensure $|U|_{0, \delta} \leq M_0$ by reducing δ . It remains, therefore, only to show that for M_1 sufficiently large and δ sufficiently small, $|v|_{1, \delta} \leq M_1$ implies $|Sv|_{1, \delta} \leq M_1$. Throughout this proof, we use M to represent any positive constant that may depend on M_1 but is otherwise independent of v and δ , and use $\tilde{M}$ for any constant that is independent of M_1, v and δ . The values of M or $\tilde{M}$ in different occurrences need not be equal.

Let $U = Sv$ and let r_i and s_i be defined by (2.5). On each branch, we show that

$$\max \{ |(r_i)_x|, |(s_i)_x| \} \leq M_1 \quad (3.2)$$

and

$$\max \{ |(r_i)_t|, |(s_i)_t| \} \leq M_1 \quad (3.3)$$

in D_δ^C , D_δ^L and D_δ^R if M_1 is large and δ is small. (Recall that D_δ^C etc. are the intersections $D_i^C \cap D_\delta$ etc., respectively.) In fact, only (3.2) needs to be shown. To see this, first observe that the vanishing initial condition and the compatibility of the initial and boundary conditions gives

$$\max_i \left\{ |P_i^B|_{C[0,\delta]}, |Q_i^B|_{C[0,\delta]} \right\} \leq M\delta.$$

Hence, we obtain from Lemma 2.1 with $\tilde{U} = 0$ that

$$|U|_{0,\delta} \leq M\delta. \quad (3.4)$$

From (2.7) and (2.9), there are constants $\tilde{M}$ and M such that

$$\begin{aligned} |\partial_i^R r_i| &\leq |l_i^R F_i| + |\partial_i^R l_i^R| |U_i| \leq \tilde{M} + M\delta, \\ |\partial_i^L s_i| &\leq |l_i^L F_i| + |\partial_i^L l_i^L| |U_i| \leq \tilde{M} + M\delta \end{aligned} \quad (3.5)$$

for each $i = 1, \dots, n$. Hence, (3.3) follows from (3.2), (3.5) and the definition of ∂_i^L and ∂_i^R in (2.8). We also note that (2.6) and (3.5) imply

$$|\partial_i^R U_i|_{0,\delta} \leq \tilde{M} + M\delta, \quad |\partial_i^L U_i|_{0,\delta} \leq \tilde{M} + M\delta \quad (3.6)$$

for all i . This will be used later.

We first consider the middle region D_δ^C , where the solution (r_i, s_i) satisfies the integral equations (2.10) and (2.12) with $r_i^I = s_i^I = 0$. Differentiating the equations with respect to x , we have

$$\begin{aligned} (r_i)_x &= (l_i^R)_x U_i(x, t) + \int_0^t [(l_i^R F_i)_x + (\partial_i^R l_i^R)(U_i)_x - (l_i^R)_x (\partial_i^R U_i)] (x_i^R)_x dt, \\ (s_i)_x &= (l_i^L)_x U_i(x, t) + \int_0^t [(l_i^L F_i)_x + (\partial_i^L l_i^L)(U_i)_x - (l_i^L)_x (\partial_i^L U_i)] (x_i^L)_x dt. \end{aligned} \quad (3.7)$$

Here, we used an identity from [6, p.469]:

$$\begin{aligned} \frac{d}{d\xi} \int_a^b f(x(t), t) Dg(x(t), t) dt \\ = f(x(b), b) g_x(x(b), b) x_\xi(b) - f(x(a), a) g_x(x(a), a) x_\xi(a) \\ + \int_a^b [f_x(x(t), t) Dg(x(t), t) - Df(x(t), t) g_x(x(t), t)] x_\xi(t) dt \end{aligned} \quad (3.8)$$

where $x(t)$ is a function such that $x(b) = \xi$ and $D = \frac{\partial}{\partial t} + x'(t) \frac{\partial}{\partial x}$. (Notice that $x_\xi(b) = 1$.) Let

$$R_i^C(t) = \sup_{\{x:(x,t) \in D_\delta^C\}} \{|(r_i)_x(x, t)|\}, \quad S_i^C(t) = \sup_{\{x:(x,t) \in D_\delta^C\}} \{|(s_i)_x(x, t)|\}. \quad (3.9)$$

From (3.4), (3.6) and (3.7), we derive

$$R_i^C(t) + S_i^C(t) \leq M\delta + M \int_0^t (1 + R_i^C(t') + S_i^C(t')) dt'$$

for $t \in [0, \delta]$. Hence, Gronwall's inequality gives

$$|(r_i)_x| \leq M\delta e^{M\delta}, \quad |(s_i)_x| \leq M\delta e^{M\delta}$$

in D_δ^C . This proves (3.2) in D_δ^C if M_1 is sufficiently large and δ is sufficiently small.

We next consider the left and right regions D_δ^L, D_δ^R which are next to an external end. Since the two cases are similar, we will consider the left region only. Let $\hat{s}_i = s_i/\varepsilon$ for any $\varepsilon > 0$. Then, the pair $(r_i, \hat{s}_i)$ satisfies the fixed point equations of either (2.14) or (2.15), depending on the type of the boundary condition. Differentiating the equations with respect to x and using a slightly modified version of (3.8) where the lower limit a of the integral also depends on ξ :

$$\begin{aligned} \frac{d}{d\xi} \int_a^b f(x(t), t) Dg(x(t), t) dt \\ = f(x(b), b) g_x(x(b), b) x_\xi(b) - f(x(a), a) g_x(x(a), a) x_\xi(a) \\ - f(x(a), a) Dg(x(a), a) a_\xi \\ + \int_a^b [f_x(x(t), t) Dg(x(t), t) - Df(x(t), t) g_x(x(t), t)] x_\xi(t) dt, \end{aligned}$$

we have

$$\begin{aligned} (r_i)_x &= (\zeta_i - l_i^R F_i - (\partial_i^R l_i^R) U_i)(0, \tau) \tau_x + (l_i^R)_x U_i(x, t) - (l_i^R)_x U_i(x_i^R)_x(0, \tau) \\ &\quad + \int_\tau^t [(l_i^R F_i)_x + (\partial_i^R l_i^R)(U_i)_x - (l_i^R)_x (\partial_i^R U_i)] (x_i^R)_x dt, \\ (\hat{s}_i)_x &= \frac{1}{\varepsilon} (l_i^L)_x U_i(t, x) + \frac{1}{\varepsilon} \int_0^t [(l_i^L F_i)_x + (\partial_i^L l_i^L)(U_i)_x - (l_i^L)_x (\partial_i^L U_i)] (x_i^L)_x dt, \end{aligned} \tag{3.10}$$

where

$$\zeta_i = 2(u_i P_i^B)_t + \varepsilon(\hat{s}_i)_t$$

if the boundary condition is given by (1.6), and

$$\zeta_i = 2 \left(\frac{a_i u_i}{\lambda_i^R} Q_i^B \right)_t + \varepsilon \left(\frac{\lambda_i^L}{\lambda_i^R} \right)_t \hat{s}_i + \varepsilon \left(\frac{\lambda_i^L}{\lambda_i^R} \right) (\hat{s}_i)_t$$

if the boundary condition is given by (1.7). This equation is valid for any ε . So, we may choose ε so small such that

$$\sigma =: \varepsilon |\lambda_i^L \tau_x(0, t)| \max \left\{ 1, \left| \left(\frac{\lambda_i^L(0, t)}{\lambda_i^R(0, t)} \right) \right| \right\} < 1, \quad t \in [0, \delta].$$

To proceed further, we need an estimate of $|\tau_x(0, t)|$. Observe that $\tau(x)$ satisfies the equation

$$x_i^R(\tau; x, t) = 0$$

where $x_i^R(\tau; x, t)$ is the solution of the initial value problem

$$\frac{dx_i^R}{ds} = \lambda_i^R(x_i^R, s), \quad x_i^R(t; x, t) = x.$$

By differentiation,

$$\lambda_i^R(0, \tau(x)) \tau_x + \frac{\partial x_i^R}{\partial x} \Big|_{(\tau(x); x, t)} = 0. \quad (3.11)$$

Let $w_i = \partial x_i^R / \partial x$. Then, w_i is the solution to the linear equation

$$\frac{dw_i}{ds} = (\lambda_i^R)_x(x_i^R(s; x, t), s) w_i, \quad w_i(t) = 1.$$

Solving the equation,

$$w_i(s) = \exp \left(\int_t^s (\lambda_i^R)_x(x_i^R(s'; x, t), s') ds' \right).$$

Returning to (3.11), we find

$$\tau_x = \frac{-1}{\lambda_i^R(0, \tau(x))} \exp \left(\int_t^{\tau(x)} (\lambda_i^R)_x(x_i^R(s'; x, t), s') ds' \right).$$

Observe that $0 < \tau(x) < t \leq \delta$ and the integrand is bounded. Hence,

$$|\tau_x| \leq \tilde{M} e^{M\delta}. \quad (3.12)$$

This is the estimate we need. By this estimate, for any M_1 , we can choose δ small enough such that the constants σ and ε are independent of M_1 . Let $R_i^L(t)$ and $\hat{S}_i^L(t)$ be defined as in (3.9) except that s_i is substituted by $\hat{s}_i$ and D_δ^C is substituted by $D_\delta^L \cup D_\delta^C$. We derive from (3.10) and the identity

$$(\hat{s}_i)_t = \partial_i^L \hat{s}_i - \lambda_i^L(\hat{s}_i)_x$$

that

$$R_i^L(t) + \hat{S}_i^L(t) \leq \sigma \hat{S}_i^L(t) + \tilde{M} + M\delta + M \int_0^t \left(1 + R_i^L(t') + \hat{S}_i^L(t') \right) dt'.$$

Replacing M and $\tilde{M}$ by $M(1 - \sigma)$ and $\tilde{M}(1 - \sigma)$, respectively, and applying Gronwall's inequality, we obtain

$$R_i^L(t) + \hat{S}_i^L(t) \leq (\tilde{M} + M\delta) e^{M\delta}.$$

Since $|s_i| \leq |\hat{s}_i|$, it follows that

$$\max \{|(r_i)_x|, |(s_i)_x|\} \leq (\tilde{M} + M\delta) e^{M\delta}$$

in $D_\delta^L \cup D_\delta^C$. This proves (3.2) in $D_\delta^L \cup D_\delta^C$ if M_1 is large and δ is small.

We next consider the case where the end of the branch is a branching or transitional junction. As before, all the branches that are connected to the same junction are considered simultaneously. Differentiating the fixed point equation $w = Kw$ where w and Kw are defined in (2.19) and (2.20), respectively, we obtain (3.10) in $D_\delta^L \cup D_\delta^C$ for $i = j_{\nu+1}, \dots, j_\mu$ and

$$\begin{aligned} (\hat{r}_i)_x &= \frac{1}{\varepsilon} (l_i^R)_x U_i(x, t) + \frac{1}{\varepsilon} \int_0^t [(l_i^R F_i)_x + (\partial_i^R l_i^R)(U_i)_x - (l_i^R)_x (\partial_i^R U_i)] (x_i^R)_x dt, \\ (s_i)_x &= (\theta_i - l_i^L F_i - (\partial_i^L l_i^L) U_i)(1, \tau) \tau_x + (l_i^L)_x U_i(x, t) - (l_i^L)_x U_i(x_i^L)_x(1, \tau) \\ &\quad + \int_\tau^t [(l_i^L F_i)_x + (\partial_i^L l_i^L)(U_i)_x - (l_i^L)_x (\partial_i^L U_i)] (x_i^L)_x dt, \end{aligned} \quad (3.13)$$

in $D_\delta^C \cup D_\delta^R$ for $i = j_1, \dots, j_\nu$, where

$$\begin{aligned} \zeta_i &= \varepsilon \sum_{l=1}^\nu (n_{j_l}^i \hat{r}_{j_l})_t(1, \tau) + \varepsilon \sum_{l'=\nu+1}^\mu (n_{j_{l'}}^i \hat{s}_{j_{l'}})_t(0, \tau) + H_i, \\ \theta_i &= \varepsilon \sum_{l=1}^\nu (m_{j_l}^i \hat{r}_{j_l})_t(1, \tau) + \varepsilon \sum_{l'=\nu+1}^\mu (m_{j_{l'}}^i \hat{s}_{j_{l'}})_t(0, \tau) + H_i, \end{aligned}$$

and m_j^i, n_j^i are defined in the proof of Theorem 2.1. Note that the estimate (3.12) holds for τ_x in both (3.10) and (3.13), although in the latter case, τ is the t -coordinate of the intersection of the left-going characteristic curve x_i^L with the vertical line $x = 1$. The derivation is identical. Hence, there is a constant ε , independent of M_1 , such that

$$\begin{aligned} \varepsilon |\tau_x| \left(\sum_{k=1}^\nu |m_{j_k}^i(t)| + \sum_{k'=\nu+1}^\mu |m_{j_{k'}}^i(t)| \right) &< 1, \\ \varepsilon |\tau_x| \left(\sum_{k=1}^\nu |n_{j_k}^i(t)| + \sum_{k'=\nu+1}^\mu |n_{j_{k'}}^i(t)| \right) &< 1 \end{aligned}$$

in $[0, \delta]$. Let σ be the maximum of the quantities on the left hand side of the above inequalities. Define $\hat{R}_i^R, S_i^R, R_i^L$ and $\hat{S}_i^L$ as in (3.9) with obvious modifications. We see that the function

$$W(t) = \sum_{l=1}^\nu \left(\hat{R}_{j_l}^R(t) + S_{j_l}^R(t) \right) + \sum_{l'=\nu+1}^\mu \left(R_{j_{l'}}^L(t) + \hat{S}_{j_{l'}}^L(t) \right)$$

satisfies the inequality

$$\begin{aligned} (1 - \sigma) W(t) &\leq \sum_{l=1}^{\nu} \left((1 - \sigma) \hat{R}_{j_l}^R(t) + S_{j_l}^R(t) \right) + \sum_{l'=\nu+1}^{\mu} \left(R_{j_{l'}}^L(t) + (1 - \sigma) \hat{S}_{j_{l'}}^L(t) \right) \\ &\leq \tilde{M} + M\delta + M \int_0^t (1 + W(t')) dt'. \end{aligned}$$

Hence, by rescaling and using Gronwall's inequality, we achieve

$$W(t) \leq (\tilde{M} + M\delta) e^{M\delta}.$$

This proves that

$$\max \{ |(r_i)_x|, |(s_i)_x| \} \leq M_1$$

in D_δ^R for $i = j_1, \dots, j_\nu$ and in D_δ^L for $i = j_{\nu+1}, \dots, j_\mu$ if M_1 is sufficiently large and δ is sufficiently small. We have thus proved (3.2) in this case.

This completes the proof of (3.2) in all cases. By choosing appropriate values of M_1 and δ , we thus obtain a set X_{δ, M_0} in the form of (3.1) which is invariant under the mapping S .

We now show that S is a contraction in X_{δ, M_0} . Let $U = Sv$, $\tilde{U} = S\tilde{v}$ for some $v, \tilde{v} \in X_\delta$, and let $W = U - \tilde{U}$. W satisfies the vanishing initial and external boundary conditions and its differential equations takes the form of (1.13) with the coefficients

$$a_i = a_i(x, t, v), \quad b_i = b_i(x, t, v), \quad c_i = c_i(x, t, v), \quad A_i = A_i(x, t, v)$$

the forcing functions f_i and g_i replaced by

$$\hat{f}_i =: f_i(x, t, v) - f_i(x, t, \tilde{v}) + (a_i(x, t, v) - a_i(x, t, \tilde{v})) \frac{\partial \tilde{Q}_i}{\partial x} \quad (3.14)$$

and

$$\begin{aligned} \hat{g}_i =: & g_i(x, t, v) - g_i(x, t, \tilde{v}) + (b_i(x, t, v) - b_i(x, t, \tilde{v})) \frac{\partial \tilde{P}_i}{\partial x} \\ & + 2(c_i(x, t, v) - c_i(x, t, \tilde{v})) \frac{\partial \tilde{Q}_i}{\partial x}, \end{aligned} \quad (3.15)$$

respectively, and the functions C_i in (2.1) replaced by

$$\begin{aligned} \hat{C}_{j_l} &= C_{j_l}(x, t, v) - C_{j_l}(x, t, \tilde{v}) + (A_{j_l}(x, t, v) - A_{j_l}(x, t, \tilde{v})) (\tilde{P}_{j_l} - \tilde{P}_{j_{unc}}), \\ \hat{C}_{j_{l'}} &= C_{j_{l'}}(x, t, v) - C_{j_{l'}}(x, t, \tilde{v}) + (A_{j_{l'}}(x, t, v) - A_{j_{l'}}(x, t, \tilde{v})) (\tilde{P}_{j_{unc}} - \tilde{P}_{j_{l'}}) \end{aligned}$$

for $l = 1, \dots, \nu$, $l' = \nu + 1, \dots, \mu$. By the Lipschitz property and the boundedness $\left| \tilde{U} \right|_{1,\delta} \leq M_1$, there is a constant M such that

$$\left| \hat{f} \right|_{0,\delta} \leq M |v - \tilde{v}|_{0,\delta}, \quad \left| \hat{g} \right|_{0,\delta} \leq M |v - \tilde{v}|_{0,\delta}, \quad \left| \hat{C} \right|_{0,\delta} \leq M |v - \tilde{v}|_{0,\delta}$$

Hence, by Theorem 2.1,

$$|Sv - S\tilde{v}|_{0,\delta} \leq M\delta |v - \tilde{v}|_{0,\delta}.$$

Therefore, S is contracting in X_{δ,M_0} if δ is sufficiently small.

The rest is standard (cf. e.g., [6]). Starting with a $v_0 \in X_{\delta,M_0}$, we generate an iterative sequence $v_{k+1} = Sv_k$. Clearly, each v_k lies in X_{δ,M_0} and the sequence converges uniformly. The limit then satisfies the integral equations in the proof of Theorem 2.1, and hence, is differentiable. Therefore, it is the solution of the quasilinear differential equations. This proves the existence and uniqueness of the solution when $U^I = 0$.

If $U^I \neq 0$, we regard U^I as a vector function of x and t and introduce $\tilde{U} = U - U^I$. It follows that $\tilde{U}$ is a solution of the quasilinear equations (1.13) with the forcing functions $\tilde{f}_i$ and $\tilde{g}_i$ given by

$$\tilde{f}_i = f_i - (Q_i^I)_x a_i, \quad \tilde{g}_i = g_i - (P_i^I)_x b_i - (Q_i^I)_x 2c_i$$

and the boundary functions are given by

$$\tilde{P}_i^B = P_i^B - P_i^I, \quad \tilde{Q}_i^B = Q_i^B - Q_i^I,$$

and

$$\tilde{C}_{j_l} = C_{j_l} + A_{j_l} P_{j_l}^I, \quad \tilde{C}_{j_{l'}} = C_{j_{l'}} - A_{j_{l'}} P_{j_{l'}}^I$$

for $l = 1, \dots, \nu$, $l' = \nu + 1, \dots, \mu$. Since $\tilde{U}$ has the vanishing initial values, it can be uniquely solved for an interval of $t \in [0, \delta]$. This gives rise to a solution U . ■

Remark: Examples can be constructed to show that if the condition (2.4) fails at $t = 0$, then, the local solution need not exist or may be not unique. In particular, if (2.4) fails at a source end, then, the system is under-determined, and if it fails at a terminal end, the system is over-determined.

We give next a result for the continuity of dependence of the solution and its derivatives on the initial, boundary and forcing functions and their derivatives. This follows from an argument similar to the proofs of Lemma 2.1 and Theorem 3.1.

Corollary 3.1 *Let $U = (P, Q)$ and $\tilde{U} = (\tilde{P}, \tilde{Q})$ be two solutions of the quasilinear problem of Theorem 3.1. Suppose the conditions of that theorem hold for the initial and boundary*

functions of both solutions. Then, there exists a constant $M > 0$, independent of initial, boundary and forcing functions, such that

$$\begin{aligned} |U - \tilde{U}|_{k,\delta} \leq M \Big(& |P^I - \tilde{P}^I|_{C^k[0,1]} + |Q^I - \tilde{Q}^I|_{C^k[0,1]} + |P^B - \tilde{P}^B|_{C^k[0,\delta]} + |Q^B - \tilde{Q}^B|_{C^k[0,\delta]} \\ & + \delta |f - \tilde{f}|_{C^k(\overline{D_\delta})} + \delta |g - \tilde{g}|_{C^k(\overline{D_\delta})} + \delta |C - \tilde{C}|_{C^k[0,\delta]} \Big). \end{aligned} \quad (3.16)$$

for $k = 0, 1$.

Proof. For $k = 0$, the result follows from substituting one of the solutions into the coefficients, modifying the forcing functions by (3.14)–(3.15), and using Lemma 2.1. For $k = 1$, we differentiate the equations and apply the lemma to the resulting equations for the derivatives of the solution. The process is standard and is omitted. ■

References

- [1] Christian Almeder, Felix Breiteneker, *et al*, Modelling of the human arterial network for an expert system for preoperative predictions, in *Proceedings of the 2nd Mathmod*, Vienna, Austria, Feb. 5-7, 1997.
- [2] Charbel, F.T., Clark, M.E., *et al*, The application of a computerized model of the cerebral circulation in skull base surgery, *2nd Intern. Skull Base Congress, San Diego*, 210, 1996.
- [3] Charbel, F.T., Misra, M., Clark, M.E., and Ausman, J.I., Computer simulation of cerebral blood flow in Moyamoya and the results of surgical therapies, *Clinical Neurology and Neurosurgery, Supp. 2*, 1997.
- [4] Clark M.E., Zhao M., Loth F., Alperin N., Sadler L., Guppy K., Charbel F.T., A patient-specific computer model for prediction of clinical outcomes in the cerebral circulation using MR flow measurements, *MICCAI*, Cambridge, England, 1999.
- [5] Clark, M.E. and Kufahl, R.H., Simulation of the cerebral macrocirculation, *Cardiovascular Systems Dynamics*, M.I.T. Press, pp. 380–390, 1978.
- [6] Courant, R. and Hilbert, D., *Methods of Mathematical Physics*, Vol. II, John Wiley & Sons, 1962.
- [7] Kufahl, R.H. and Clark, M.E., A circle of Willis simulation using distensible vessels and pulsatile flow, *J. Biomechanical Engineering* 107, pp. 112-122, 1985.

- [8] McOwen R., *Partial Differential Equations*, Prentice Hall, 1995.
- [9] Porenta, G., Young, D.F., Rogge, T.R., A finite-element model of blood flow in arteries including taper, branches, and obstructions, *J. Biomechanical Engineering*, Vol. 108, pp. 161-167, 1986.
- [10] Raines, Jeffrey K., Yaffrin, Michel Y. and Shapiro, Ascher H., A computer simulation of the human arterial system, *Proceedings of the 1971 Summer Computer Conference*, Vol. 2, pp. 171-178.
- [11] Raines, Jeffrey K., Yaffrin, Michel Y. and Shapiro, Ascher H., A computer simulation of arterial dynamics in the human leg, *J. Biomechanics*, Vol. 7, pp. 77-91, 1974.
- [12] Ruan, Weihua, Clark, M.E., Zhao, Meide and Curcio, Anthony, A hyperbolic system in a one-dimensional network, arXiv: math-ph/0209015.
- [13] Smith, N.P., Pullan, A.J. and Hunter, P.J., An anatomically based model of transient coronary blood flow in the heart, *SIAM J. Appl. Math.*, Vol. 62, pp. 990–1018, 2002.
- [14] Spaan, J.A., Breuls, N.P.W. and Laird, J.D., Diastolic-systolic coronary flow differences are caused by intramyocardial pump action in the anesthetised dog, *Circ. Res.* Vol. 49, pp. 584-593, 1981.
- [15] Zhao M., Charbel F.T., Alperin N., Loth F., Clark M.E., Improved phase-contrast flow quantification by three-dimensional vessel localization, *Mag. Reson. Imag.*, Vol. 18, 697-706, 2000